

**ECULIZUMAB (SOLIRIS®)
PRESCRIBER ORDER FORM**



Fax completed form, insurance information, and clinical documentation to:

Warren, MI t 888-386-0886 f 586-263-3306

Patient Name:	Date of Birth:
---------------	----------------

Address:

Phone:	Height:	<input type="checkbox"/> inches <input type="checkbox"/> cm	Weight:	<input type="checkbox"/> lbs <input type="checkbox"/> kg
--------	---------	---	---------	--

CLINICAL INFORMATION

Primary Diagnosis Description:	ICD-10 Code:
--------------------------------	--------------

Meningococcal Vaccination Status:

Primary vaccination series completed – date: _____

MenACWY booster completed – date: _____

MenB booster completed – date: _____

ECULIZUMAB (SOLIRIS®) PRESCRIPTION

Eculizumab (Soliris®) refill as directed x 1 year

Induction Dose: Infuse 600 mg IV over at least 35 min weekly x 4 weeks.
 Infuse 900 mg IV over at least 35 min weekly x 4 weeks.
 Other: _____

Maintenance Dose: Infuse 900 mg IV over at least 35 min on Week 5, then every 2 weeks thereafter.
 Infuse 1200 mg IV over at least 35 min on Week 5, then every 2 weeks thereafter.
 Infuse _____ mg IV over at least 35 min every 2 weeks.
 Other: _____

Max infusion time not to exceed 2 hours.

ANCILLARY ORDERS

Anaphylaxis Kit

If this is a 1st dose, would you like Compassus to provide an anaphylaxis kit with the 1st dose?

Yes No

- Dosage:**
- Epinephrine 0.3 mg (> 30 kg), 0.15 mg (15 to 30 kg), or 0.01 mg/kg (< 15 kg) SQ or IM x 1; repeat x 1 in 5 to 15 min PRN.
 - Diphenhydramine 25mg PO or IV/IM. May repeat additionally 25mg PO or IV PRN.
 - Normal saline 500 mL (> 30 kg) or 250 mL (< 30 kg) IV at KVO rate PRN anaphylaxis. Patient < 30 kg, infuse over 2 to 4 hours PRN headache rated > 5 on pain scale.

Medication Orders

Acetaminophen 650 mg PO 30 min before infusion. Patient may decline.
 Diphenhydramine 25 mg PO 30 min before infusion. Patient may decline.
 Other: _____

IV Flush Orders

Peripheral: NS 2 to 3 mL pre-/post-use.
 Implanted Port: NS 5 to 10 mL pre-/post-use and 10 to 20 mL pre-/post-lab draw. Heparin (100 unit/mL) 3 to 5 mL post-use. For maintenance, heparin (100 unit/mL) 3 to 5 mL every 24 hr if accessed or weekly to monthly if not accessed.

Lab Orders

No labs ordered at this time.
 Other: _____

Skilled nurse to initiate IV access for administration of doses in the home or alternate care setting. Access to be discontinued upon completion of infusion. Refill above ancillary orders as directed x 1 year.

I certify that the use of the indicated treatment is medically necessary and I will be supervising the patient's treatment.

Prescriber Signature: _____ **Date:** _____

PRESCRIBER INFORMATION

Prescriber Name:	Phone:	Fax:
------------------	--------	------

Address:	NPI:
----------	------

City, State:	Zip:	Office Contact:
--------------	------	-----------------

CONFIDENTIAL HEALTH INFORMATION: Healthcare information is personal information related to a person's healthcare. It is being faxed to you after appropriate authorization or under circumstances that don't require authorization. You are obligated to maintain it in a safe, secure, and confidential manner. Re-disclosure of this information is prohibited unless permitted by law or appropriate customer/patient authorization is obtained. Unauthorized re-disclosure or failure to maintain confidentiality could subject you to penalties described in federal and state laws. **IMPORTANT WARNING:** This message is intended for the use of the person or entity to whom it is addressed and may contain information that is privileged and confidential, the disclosure of which is governed by applicable law. If the reader of this message is not the intended recipient, or the employee or agent responsible for delivering it to the intended recipient, you are hereby notified that any dissemination, distribution or copying of this information is STRICTLY PROHIBITED. If you have received this message in error, please notify us immediately. Brand names are the property of their respective owners.