



IMMUNOGLOBULIN (IG)
IV AND SUBQ ORDERS

P: 888-386-0886 | F: 586-263-3306

PATIENT INFORMATION:

Fax completed form, insurance information and clinical documentation to 586-263-3306

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_

Last 4 of SS #: \_\_\_\_\_ Patient Status: [ ] New [ ] Continuing Therapy Next Treatment Date: \_\_\_\_\_

MEDICAL INFORMATION

ICD-10 Code (required): \_\_\_\_\_ ICD-10 descriptions: \_\_\_\_\_

Patient weight: \_\_\_\_\_ kg Height: \_\_\_\_\_ Diabetic: [ ] Yes [ ] No If obese, use adjusted body wt? [ ] Yes [ ] No

Allergies: \_\_\_\_\_ Brand previously used: \_\_\_\_\_

THERAPY ORDER

[ ] IV [ ] SQ Pharmacist to identify clinically appropriate brand/infusion rates. May substitute based on product availability.

Table with 4 columns: Dose Type, Units, Frequency, and Administration. Rows for Loading Dose and Maintenance Dose.

[ ] Do not substitute. Administer brand: \_\_\_\_\_

- Infuse entire contents of Ig infusion bag/vial(s) per current dose.
If needed, round dose to nearest whole 5 gm vial for IV doses and nearest single-use vial size for subQ doses.

Premedication orders: to be administered 15-30 minutes before infusion

- Acetaminophen 500mg PO, Solu-Medrol mg IVP, Loratadine 10mg PO, Normal Saline 500mL IV, Diphenhydramine 25mg PO, Cetirizine 10mg PO, Other:

Lab orders: \_\_\_\_\_ Lab frequency: [ ] Each infusion [ ] Other: \_\_\_\_\_

Anaphylactic Reaction Orders:

- Epinephrine (based on patient weight)
>30kg (>66lbs): EpiPen® 0.3mg or compounded syringe IM or SQ; may repeat in 5-10 minutes x 1
15-30kg (33-66lbs): EpiPen® 0.15mg or compounded syringe IM or SQ; may repeat in 5-10 minutes x 1
Diphenhydramine - Administer 25-50mg orally OR IV (adult), refer to provider orders or policy for pediatric dose
NS 500 mL IV bolus as needed for IVIg therapy (adult), refer to provider orders or policy for pediatric bolus

Flush orders: NS 1-20mL pre/post infusion PRN and Heparin 10U/mL or 100U/mL per protocol as indicated PRN

\*FOR COMPASSUS USE ONLY

Drug/Brand Selection: \_\_\_\_\_ Date: \_\_\_\_\_

NP/Pharmacist Name: \_\_\_\_\_ NP/Pharmacist Signature: \_\_\_\_\_

PROVIDER INFORMATION

By signing this form and utilizing our services, you are authorizing Compassus and its employees to serve as your prior authorization and specialty pharmacy designated agent in dealing with medical and prescription insurance companies, and to select the preferred site of care for the patient.

Provider Name: \_\_\_\_\_ Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Provider NPI: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Contact Person: \_\_\_\_\_

[ ] Opt out of Compassus selecting site of care (if checked, please list site of care) \_\_\_\_\_

PREFERRED LOCATION

City: \_\_\_\_\_ State: \_\_\_\_\_

Compassus

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee and contains material that is confidential, privileged property, or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately and destroy all copies if you have received this document in error.

**REQUIRED DOCUMENTATION FOR INSURANCE APPROVAL**  
GENERAL REQUIREMENTS

- Patient demographics
- Insurance information
- All applicable diagnoses
- History and physical
- Recent progress notes within 12 months
- Patient's height and weight
- Drug allergies
- Physician Orders
- Plus one of the following

COMMON VARIABLE IMMUNODEFICIENCY (CVID) /  
HYPOGAMMAGLOBULINEMIA / PARKINSON'S DISEASE (PD)

- Lab last showing Ig levels and subclasses Ig levels.
- Documentation of recurrent infections
- History of antibiotic usage - showing failure to respond to antibiotics
- Documented inadequate response to pneumococcal vaccine or tetanus/diphtheria

CHRONIC INFLAMMATORY DEMYELINATING POLYNEUROPATHY (CIDP) /  
GUILLAIN-BARRÉ SYNDROME (GBS)

- Labs
- Nerve conduction study, electromyography (EMG)
- Nerve and/or muscle biopsy (if available)
- Nerve conduction velocity (NCV) test results
- Tried and failed treatments
- Spinal tap (if available)

MYASTHENIA GRAVIS

- Exacerbation
- Any history of crisis
- Thymectomy
- Any symptoms that affect respiration, speech or motor function
- Tried and failed treatments